

WNL = within normal limits

PRE-SALE EQUINE PHYSICAL SURVEY

Horse's Name Crescentzee Breed Thoroughbred
Year Foaled '17 Color Gray Sex Filly

Consignor - First Daniel Last Kerr Owner Agent
Owner - First Daniel & Angela Last Kerr

REMINDER - ORIGINAL NEGATIVE COGGINS DRAWN WITHIN 12 MONTHS OF THE DATE OF THE AUCTION REQUIRED

Date of Examination: 5/15/20 Place of Examination: Kerr Racing Horses Keizer, OR

Medical History (Colic, Founder, Surgery, Intestinal Disorders, Lameness, Respiratory, Nerves, etc.)

leg wounds from fence injury & trailer injury. was ultrasounded and found no damage to flexor tendons.

Clinical Evaluation

Body Temperature: 99.0 Eyes: WNL Mouth: WNL
Skin: WNL Tumors: none Scars: must client to lower neck legs

Cardiovascular (Heart Rate /Respiratory): WNL

Evidence of Bleeder: none Gastrointestinal / Feces: normal

Neurological / Musculoskeletal: WNL Heart Rate 36 Resp rate 12

Equine Physical Exam

Indication of Lameness: none Evidence of Founder or Laminitis: none

Feet: Left Fore: WNL Right Fore: WNL
Left Hind: WNL Right Hind: WNL

Limbs (Examine for lameness, enlargements, abnormalities) - all 4 legs conformation WNL

Left Fore: scar dorsal metacarpus Right Fore: WNL
Left Hind: scar over lower flexor tendon Right Hind: scar mid flexor tendon

Urogenital (Penis, Testicles, Prepuce) (Vulva, Vagina, Urethral Orifice, Cervix): WNL externally

Broodmares - Vaginal Exam: Culture (on open mare being offered as broodmare): _____

Broodmares - Pregnant: _____ If open - Palpation (Ovaries, Uterus, Cervix): _____

Comments, Observations and Recommendations: _____

Examining Veterinarian: Michael J. JDM Date: 5/15/20

Address: P.O. Box 409 Silverton, OR 97381

Phone: 503-967-4404

Flashpoint Bloodstock, LLC has the right to refuse any horse based on the results of this examination or any information known to the consignee regarding suitability of sale. If there is any doubt as to the suitability of any horse for sale purposes, notify the sale management.



Client Details

Name Kerr, Daniel
Address 2828 Zee Lane Northeast
Salem, Oregon, 97303

Phone(s) Mobile: 971-304-5258

Patient Details

Name Crescent
Species Equine (Horse)
Breed Thoroughbred
Age 2 years 2
Sex Filly

10-14-2019

Laceration Discharge Summary for Crescent

Case Summary: Crescent, 2yo Thoroughbred mare, presented into LBEMC for a right hind metatarsal ultrasound following laceration to the plantar metatarsal region. Last Friday the wound was examined 24 hours post-injury and ultrasound was recommended due to concern for tendon involvement. The wound could not be sutured due to extensive limb swelling. Upon arrival at LBEMC Crescent was found to have sustained a laceration to her right forehead during the trailer ride. Reported to be up-to-date on tetanus vaccine within the past 6 months.

Temperature: 100.2 F

Heart Rate: 48 bpm

Respiration: 20 brth/min

Attitude: Bright, alert, and responsive

Lameness: None at the walk

Laceration (head): There is a fresh full-thickness 'v' shaped laceration on the right forehead. The wound is approximately 6-7cm in length.

Laceration (right hind): Bandage in place upon arrival. Upon removal of the bandage, there is a significant improvement in limb swelling. There is a healthy bed of granulation tissue in the wound and the wound flap is moderately contracted. No foul smell or increased wound drainage.

Right hind metatarsal ultrasound: There was thickening of the area around the wound with gas near the bottom of the wound. The tendons (superficial and deep digital flexor tendons) did not appear to be affected by the laceration. There was no obvious fracture of the splint bone on the outside of the limb. As discussed this tendon is at risk forming adhesions due to the close proximity of the wound to the tendon. This wound is likely going to take 2-3 months to heal completely.

The laceration on the forehead was clipped, cleaned, blocked locally, and sutured closed using 1 PDS in a simple interrupted pattern.

Medications

1. Trimethoprim-Sulfa (TMS antibiotic): Please give 16 tabs orally twice daily for 14 days.
2. Discontinue penicillin and gentamicin.
3. Bute: Please give 1 gram orally twice daily for 2 days, then 1 gram orally once daily for 2 more days.

*Many medications can cause diarrhea in horses. If your horse develops loose manure or stops eating, please discontinue all medications and call us right away.

Exercise/Housing

Stall confinement until the wound is healed.

Bandage Changes

Please change the bandage every 2-3 days. It is important that she remains in a heavy compression bandage until the wound is healed in order to reduce motion, proud flesh, and swelling. Bandage layers are as follows:

- 1) Neosporin antibiotic ointment on wound (may need to use steroid ointment if proud flesh develops). Non-adherent pad held in place with gauze roll.
- 2) Thick cotton padding, brown gauze, vet wrap. Be sure the brown gauze and vet wrap does not contact the skin. Elastikon at top and bottom.

Recheck Exam: Upon Crescent arriving back home please have your regular vet re-examine and take over her care.

Suture Removal: Please have her sutures removed in 14-18 days

Don't hesitate to contact us with any questions or concerns.

Rudy Madrigal DVM, DACVIM
Grace Monmaney DVM,
gmonmaney@lbemc.com
and the team at Loomis Basin Equine Medical Center



Client Details

Name Kerr, Daniel
Address 2828 Zee Lane Northeast
Salem, Oregon, 97303

Phone(s) Mobile: 971-304-5258

Patient Details

Name Crescent
Species Equine (Horse)
Breed Thoroughbred
Age 2 years 2
Sex Filly

10-14-2019

Laceration Discharge Summary for Crescent

Case Summary: Crescent, 2yo Thoroughbred mare, presented into LBEMC for a right hind metatarsal ultrasound following laceration to the plantar metatarsal region. Last Friday the wound was examined 24 hours post-injury and ultrasound was recommended due to concern for tendon involvement. The wound could not be sutured due to extensive limb swelling. Upon arrival at LBEMC Crescent was found to have sustained a laceration to her right forehead during the trailer ride. Reported to be up-to-date on tetanus vaccine within the past 6 months.

Temperature: 100.2 F

Heart Rate: 48 bpm

Respiration: 20 brth/min

Attitude: Bright, alert, and responsive

Lameness: None at the walk

Laceration (head): There is a fresh full-thickness 'v' shaped laceration on the right forehead. The wound is approximately 6-7cm in length.

Laceration (right hind): Bandage in place upon arrival. Upon removal of the bandage, there is a significant improvement in limb swelling. There is a healthy bed of granulation tissue in the wound and the wound flap is moderately contracted. No foul smell or increased wound drainage.

Right hind metatarsal ultrasound: There was thickening of the area around the wound with gas near the bottom of the wound. The tendons (superficial and deep digital flexor tendons) did not appear to be affected by the laceration. There was no obvious fracture of the splint bone on the outside of the limb. As discussed this tendon is at risk forming adhesions due to the close proximity of the wound to the tendon. This wound is likely going to take 2-3 months to heal completely.

The laceration on the forehead was clipped, cleaned, blocked locally, and sutured closed using 1 PDS in a simple interrupted pattern.

Medications

1. Trimethoprim-Sulfa (TMS antibiotic): Please give 16 tabs orally twice daily for 14 days.
2. Discontinue penicillin and gentamicin.
3. Bute: Please give 1 gram orally twice daily for 2 days, then 1 gram orally once daily for 2 more days.

*Many medications can cause diarrhea in horses. If your horse develops loose manure or stops eating, please discontinue all medications and call us right away.

Exercise/Housing

Stall confinement until the wound is healed.

Bandage Changes

Please change the bandage every 2-3 days. It is important that she remains in a heavy compression bandage until the wound is healed in order to reduce motion, proud flesh, and swelling. Bandage layers are as follows:

- 1) Neosporin antibiotic ointment on wound (may need to use steroid ointment if proud flesh develops). Non-adherent pad held in place with gauze roll.
- 2) Thick cotton padding, brown gauze, vet wrap. Be sure the brown gauze and vet wrap does not contact the skin. Elastikon at top and bottom.

Recheck Exam: Upon Crescent arriving back home please have your regular vet re-examine and take over her care.

Suture Removal: Please have her sutures removed in 14-18 days

Don't hesitate to contact us with any questions or concerns.

Rudy Madrigal DVM, DACVIM
Grace Monmaney DVM,
gmonmaney@lbemc.com
and the team at Loomis Basin Equine Medical Center